

## Original Article

**Cite this article:** Lodi L, Maricato G (2020). Rough-toothed dolphins (Cetartiodactyla: Delphinidae) habitat use in coastal urban waters of the South-western Atlantic. *Journal of the Marine Biological Association of the United Kingdom* 1–9. <https://doi.org/10.1017/S0025315420000132>

Received: 5 November 2018

Revised: 16 January 2020

Accepted: 17 January 2020

### Key words:

Cetacean; group size and composition; habitat use; movements; photo-identification; Rio de Janeiro; seasonality; site fidelity; *Steno bredanensis*

### Author for correspondence:

Liliane Lodi, E-mail: [lilianelodi@gmail.com](mailto:lilianelodi@gmail.com)

# Rough-toothed dolphins (Cetartiodactyla: Delphinidae) habitat use in coastal urban waters of the South-western Atlantic

Liliane Lodi<sup>1</sup>  and Guilherme Maricato<sup>1,2</sup> 

<sup>1</sup>Projeto Ilhas do Rio and Projeto Baleias & Golfinhos do Rio de Janeiro, Instituto Mar Adentro, Rua Siqueira Campos 33, sala 504, Copacabana, Rio de Janeiro, RJ, CEP 22031-071, Brazil and <sup>2</sup>Laboratório de Bioacústica e Ecologia de Cetáceos, Universidade Federal Rural do Rio de Janeiro, Rodovia BR 465, km 7, Seropédica, Rio de Janeiro, RJ, CEP 23897-000, Brazil

## Abstract

Rough-toothed dolphins (*Steno bredanensis*) are regularly present in continental shelf areas of the South-western Atlantic. However, there is little information on the natural history and ecology of these delphinids. This study evaluated the occurrence, habitat use and individual movements of the species in coastal waters off Rio de Janeiro, south-eastern Brazil. Data were obtained from boat surveys between August 2011 and May 2018, during which rough-toothed dolphins were sighted in 21 distinct events, predominantly in autumn and winter. The mean group size was 29 individuals. Rough-toothed dolphins were usually recorded 130 to 2300 m from the coast, between 7.6 and 28 m depths. In total, 115 individuals were catalogued through dorsal fin marks and 61 (53%) were resighted between one (47.5%) and four (9.8%) occasions. The interval between resightings ranged from seven to 2087 days (mean = 268). Agglomerative hierarchical clustering indicated 30 individuals (49.2%) in low degree, 12 (19.7%) in medium degree and 19 (31.1%) in high degree of site fidelity. Dolphins showed a higher frequency of low degree of habitat use, despite the presence of multiyear recaptures, which may be related to the prevalence of dolphin occurrence in autumn and winter, a large home range and/or the abundance and distribution of food resources. Dedicated surveys and regional collaboration are needed to evaluate the home range and population status of this species for their effective conservation. Our findings enhanced knowledge of this little studied species facing increasing anthropogenic threats in coastal waters off Rio de Janeiro.

## Introduction

Although not commonly observed in specific sites in the South-western Atlantic, rough-toothed dolphins (*Steno bredanensis*) (Lesson, 1828) are widely distributed over the continental shelf (Lodi & Hetzel, 1998b; Rossi-Santos *et al.*, 2006) but little information is available on their occurrence outside coastal areas. Few records of the species in deep oceanic waters (1500 and 4123 m) are available (Ramos *et al.*, 2010; Wedekin *et al.*, 2014).

Distribution results report pelagic habitat usage of this species, for most part of its range, between 1000 to 2000 m depths off French Polynesia (Gannier & West, 2005) and greater than 1500 m depths off Hawaii (Baird *et al.*, 2008). In Rio de Janeiro, south-eastern Brazil, the rough-toothed dolphin occurs closer to the coast. They are most commonly found at depths of 5–20 m (Lodi & Tardin, 2018).


Their offshore distribution in most tropical and subtropical areas of all three major oceans should minimize potential threats arising from habitat loss and/or habitat alteration, whereas such threats are reported for coastal areas, where potential impacts from human activities would be higher. Rough-toothed dolphin in Brazilian waters may be affected negatively by interactions with fisheries (Lodi & Capistrano, 1990; Di Benedetto *et al.*, 1998; Lodi & Hetzel, 1998b; Monteiro-Neto *et al.*, 2000; Netto & Barbosa, 2003), habitat degradation and chemical pollution (Dorneles *et al.*, 2007; Yogui *et al.*, 2010; Lailson-Brito *et al.*, 2012) and marine debris ingestion (Meirelles & Barros, 2007), but the magnitude of such effects remains unknown.

Studies based on individual identification of rough-toothed dolphins have been conducted around Atlantic and Pacific oceanic islands and have shown some degree of site fidelity with individuals being seen exclusively around one island in multiple years (Mayr & Ritter, 2005; Kuczaj & Yeater, 2007; Baird *et al.*, 2008; Oremus *et al.*, 2012).

Information on the individual and spatial behaviour of *S. bredanensis* in the South-western Atlantic are scarce. Individual movements between areas of south-eastern Brazil suggest that the species can possibly display site fidelity also in coastal environments (Lodi *et al.*, 2012; Wedekin *et al.*, 2017; Santos *et al.*, 2019).

While they are widely distributed throughout tropical and warm-temperate waters worldwide, little is known about rough-toothed dolphins anywhere in their range. Detailed, long-term behavioural ecology studies on rough-toothed dolphins have only been conducted in recent years and in deep waters around oceanic islands, hence our knowledge is limited to specific areas. The objective of this study was to evaluate the occurrence, habitat use and individual movements of the species in coastal waters off Rio de Janeiro, which are subject to intense





**Fig. 1.** Study area along the coast of Rio de Janeiro state, south-eastern Brazil. Isobaths are shown in continuous light grey lines. Study area (dashed line) was delimited as the polygon around survey routes, which also corresponds to the area surveyed by boat. The dashed outlines represent the MPAs (MoNa Cagarras and Resex Marinha Itaipu). Records of rough-toothed dolphin (*Steno bredanensis*) recorded during surveys along the coastal waters off Rio de Janeiro from August 2011 to April 2018.

human use, to understand their behavioural ecology and provide baseline information vital to support conservation measures.

Guanabara Bay, a eutrophic coastal bay in south-eastern Brazil, is part of a large ecosystem that forms the Guanabara Bay drainage basin, impacted by the polluted discharge from the Rio de Janeiro metropolitan area. After decades of continuous anthropogenic impact and degradation, and intense eutrophication, Guanabara Bay faces a complex of environmental, social and economic challenges (see Fistarol *et al.*, 2015 for a review). The coast of Rio de Janeiro is vulnerable to pollution from sources such as the Ipanema submarine sewer outfall, and various waste discharges in the waters of Guanabara Bay (van Weerelt *et al.*, 2013).

## Materials and methods

### Study area

Our study comprised coastal waters off the city of Rio de Janeiro, located on the eastern coast of Brazil (Figure 1) and one of the most densely populated cities in Latin America with ~6,600,000 inhabitants (IBGE, 2018).

Located to the north-east coast of the city, Guanabara Bay, historically known as a highly contaminated area (Fistarol *et al.*, 2015), is the second largest coastal bay in Brazil with an area of 384 km<sup>2</sup> (Kjerfve *et al.*, 1997). Most of the bay (84%) is <10 m depth, with a maximum depth of 58 m on its central channel (Ruellan, 1944). Tides in the Guanabara Bay are mainly semidiurnal, with a mean tidal range of ~0.7 m (spring tidal range: 1.1 m, neap tidal range: 0.3 m), without significant spatial variance (Fistarol *et al.*, 2015).

Two Marine Protected Areas (MPA) are included in the study area (Figure 1). On 13 April 2010, the Cagarras Archipelago Natural Monument (MoNaCa), a category III (Day *et al.*, 2012) was established to protect natural features and their associated biodiversity and habitats that are highly valuable for visitors and where recreational activities are encouraged. Extractive use (of living or dead material) is not deemed consistent with the objectives of this category. The Itaipu Marine Extractive Reserve (Resex Itaipu) is a category V (Day *et al.*, 2012), and it was established on 30 September 2013 by Rio de Janeiro state law number 44,417. This category applies to areas where local communities, interacting with nature over time, have produced areas of distinct character with traditional management practices such as fisheries and sustainably use the sea. Nevertheless, the primary objectives of these areas remain nature conservation.

### Survey effort

Data on rough-toothed dolphins reported here were collected between 2011 and 2018. Survey effort was non-systematic and track lines generally intended to maximize the likelihood of finding cetaceans.

### Group size and composition

Immature individuals were considered to be up to two-thirds the length of adults, classified as such through visual estimation. For the purpose of this study, immatures included neonates, calves and juveniles, and group refers to all dolphins sampled in a single

encounter, thus defined as a spatial aggregation that appears to be involved in a similar activity (e.g. foraging, socializing, resting or travelling, Shane *et al.*, 1986). Group locations were determined using a Garmin Etrex 30x Global Positioning System (GPS).

### Habitat use

Occurrence, distance from coast, depth and slope were mapped using ArcGIS Desktop 10.6. Distance from coast was considered as the smallest possible distance between each sighting and the coast. The study area was divided into a 1 × 1 km grid with 226 cells. Depths were obtained with a bathymetric raster surface (Nautical Chart no. 1506, Directorate of Hydrography and Navigation of the Brazilian Navy) using the depth closest to the dolphins' recorded location. Whenever more than one value was available, these were averaged to assign the depth of the sighting. The slope was calculated as the difference between maximum and minimum depth values in each grid divided by the distance in metres between both depth values. Then the values were converted to degrees.

First, a Multiple correspondence analysis (MCA) (Greenacre, 1988) was performed to visually represent the relations among similar variables. Moreover, quantitative variables (group size, depth, slope and distance from coast) were divided into three categories by the natural breaks using the conversion of values to factors (Supplementary Figure S1). Univariate tests were later performed to test the relation between these variables.

Seasons were defined as summer (December to February), autumn (March to May), winter (June to August) and spring (September to November). The ANOVA one-way test (DeGroot & Schervish, 2012) was used to verify significant differences in group size in different months, seasons and years sampled as well.

Analysis of group size variation in the presence of immatures was carried out with the Shapiro–Wilk normality test (Mason *et al.*, 2003). The Bartlett's test (Mason *et al.*, 2003) was then performed to determine if parameters varied independently and, finally, significance was tested with the Student's *t*-test for independent samples (DeGroot & Schervish, 2012). The same analyses were run to investigate possible temporal-spatial relations using the closing season for the fisheries of Lebranche mullet, *Mugil liza* (when mullets approach the coast to spawn in shallower waters) and the depth associated with dolphin sightings. The Lebranche mullet closing season runs from 15 March to 15 September (Ordinance No. 04, of 14 May 2015, Ministry of Fisheries and Aquaculture and the Environment), in order to ensure that the reproductive cycle of the species and their stocks are maintained.

The Fisher's exact test (Choi *et al.*, 2015) was applied for further analysis of the presence of immatures in groups sighted in relation to the type of coast (visually determined as rocky or sandy) and the closing season for Lebranche mullet fisheries. This same test was also carried out to investigate if the type of coast preferentially used by rough-toothed dolphins showed any relation to the closing season for Lebranche mullet fisheries. The choice in the use of the Fisher's *s* test was due to the existence of frequencies smaller than 5.

Tidal state data (ebb tide and flood tide) were obtained from the Centre of Hydrography of the Brazilian Navy at the nearest port of the study area (Ilha Fiscal: 22°53'8"S 43°10'0"W). The terms used to describe direction of dolphins' movements were 'up' for inwards movements towards Guanabara Bay and 'down' for outwards movements towards the sea. Thus, the Wilcoxon signed-rank test (DeGroot & Schervish, 2012) was used to analyse the movement of 'up' and 'down' of dolphins' groups in Guanabara Bay. This test investigated whether the proportion of 'up' and 'down' movements varied at different depths and slopes for the recorded sightings. To test whether movement and tide recorded for sightings were dependent, the  $\chi^2$  test (DeGroot &

Schervish, 2012) was carried out. All tests were performed using a 0.05 significance level.

### Photo-identification

Photo-identification has been established as a powerful tool in cetacean research (Whitehead *et al.*, 2000). The nicks and notches along the trailing edge of the dorsal fin define an individual's unique signature (Würsig & Jefferson, 1990). Rough-toothed dolphins exhibit distinct features suitable for individual identification, such as notch patterns on the dorsal fin and distinct scratches (Mayr & Ritter, 2005). During each encounter, an attempt was made to photograph as many individuals as possible, regardless of distinctive marks or vicinity to the vessel. Only well-focused images were considered for the analysis and creation of a photo-identification catalogue; these were unobscured, with dorsal fin perpendicular to the plane of the photographer, with the dorsal fin large enough to identify notches, and which showed the entire anterior and trailing edge of the dorsal fin from the tip to anterior and posterior insertion. The protocol for photo-identification in Espécie *et al.* (2010) were followed. One researcher was designated to focus on obtaining suitable photographs of individuals during sightings, using a digital SLR camera with 100–300 mm zoom lenses for photo-identification purposes.

All dolphin sightings and resightings from photographs were confirmed by two independent researchers before being entered into our database. Photographed individuals were classified according to their quality ratings on a scale of 1 to 5 (poor to excellent) (Oremus *et al.*, 2007) and sighting histories. Only photographs classified in scales 4 and 5 were used. To avoid mis-identification, immatures and individuals without distinctive markings were not included in the analysis. The photographs of distinct dorsal fins were compared among individuals using Darwin software 2.22 (Digital Analysis and Recognition of Whale Images on a Network: darwin.eckerd.edu).

In addition, reference catalogues were constructed with photographic records to explore individual movements and site fidelity to the area: the latter was defined as the tendency of an individual to return to an area previously occupied or remain in an area over an extended period (White & Garrot, 1990).


Site fidelity was classified in three degrees (low, medium and high) based on analysis of three parameters: (1) the ratio between the number of sightings and the number of survey dates from its first sighting to its last; (2) the ratio between the number of survey dates a dolphin was sighted and the total of survey dates; and (3) the ratio between the number of seasons a dolphin was sighted and the total of seasons (adapted from Passadore *et al.*, 2018 using only individual recaptures). We used Agglomerative hierarchical clustering (AHC) analysis to separate individuals into three categories based on the dissimilarity of these three standardized parameters (Legendre & Legendre, 2012), by means of the average distance method and the Euclidean distance. Finally, the Correlation Cophenetic Coefficient (CCC) value was calculated (Sokal & Rohlf, 1962) to ascertain if the clustering was efficient, when the result was above 0.7 (Rohlf, 1970).

All computational procedures were performed using the RStudio software 1.1.456 (<https://www.rstudio.com>) through the R 3.5.1 (<https://www.r-project.org>).

## Results

### Survey effort

Surveys were undertaken on 183 vessel days, covering 9416.8 km of trackline in 1314.55 h of effort between August 2011 and May 2018 (Table 1), during which rough-toothed dolphins were sighted in 21 distinct events (see Figure 1).



**Fig. 2.** Seasonal (A) and monthly (B) distribution of rough-toothed dolphins (*Steno bredanensis*) groups along coastal waters off Rio de Janeiro (N = 21) between August 2011 and April 2018. With records (light grey) and without records (dark grey).

**Table 1.** Survey effort and record rate of rough-toothed dolphins in coastal waters off Rio de Janeiro per year

Year	Month/Vessel surveys	Mean effort (h)	Overall effort (h)	Overall effort (km)	Record rate <sup>a</sup>
2011	Aug–Dec/20	7.8 (4.3–9.0)	157.5	1010.9	0.1
2012	Aug–Dec/21	5.8 (4.3–9.4)	122.3	1039.6	0.04
2014	Jan–Dec/35	7.3 (5.3–10.0)	257.5	1868.4	–
2015	Jan–Dec/25	7.3 (5.1–9.1)	184.4	1456.8	0.04
2016	Jan–Dec/31	6.2 (5.1–8.5)	208.5	1714.9	0.12
2017	Jan–Dec/38	7.0 (5.4–8.3)	293.3	2007.6	0.31
2018	Feb–May/13	7.0 (6.2–8.1)	91.05	318.6	0.07
Total	183		1314.55	9416.8	

Values in parentheses represent the minimum and maximum effort/year.

<sup>a</sup>Calculated as number of total groups sighted that year/number of survey effort days that year.

The group size ranged from a minimum of four to a maximum of 60 individuals (mean = 29, SD = 14). There were no significant differences in group size by month (one-way ANOVA,  $P = 0.36$ ), season (one-way ANOVA,  $P = 0.29$ ) or year (one-way ANOVA,  $P = 0.41$ ).

### Group size and composition

The presence of immature individuals in 76.2% of the sighted groups resulted in higher numbers of dolphins per group (Student's  $t$ -test,  $P = 0.03$ ).

Group records were made throughout the year with two peaks, in autumn (23.7%) and winter (17.9%) (Figure 2A). The frequency of group records per month (Figure 2B) varied from 0 to 33.3 (mean = 13.2, SD = 12.6,  $N = 12$ ).

### Habitat use

*Steno bredanensis* groups were usually recorded between 130 and 2300 m from the coast (mean = 760, SD = 545) and depths between 7.6 and 28 m (mean = 14.9, SD = 5.4). The occurrence grids showed slopes between 0 and 4.2° (mean = 0.7°, SD = 1.1°).

Groups with immatures did not show dependence on the type of coast or the closing season for Lebranche mullet fisheries (Fisher's exact test and  $P = 0.22$  and 0.55, respectively). However, the type of coast where groups were sighted has shown to be related to the closing season for Lebranche mullet fisheries (Fisher's exact test,  $P = 0.02$ ). Groups were sighted with higher frequency near rocky shores during the closing season for Lebranche mullet fisheries.

Depth and slope recorded at dolphin sighting locations were not significantly different during the closing season for Lebranche mullet fisheries (Student's  $t$ -test,  $P = 0.1761$  and Wilcoxon test,  $P = 0.0904$ , respectively).

The movement of rough-toothed dolphin groups in Guanabara Bay was related to depth, with more sightings 'up' the bay at greater depths (Wilcoxon test,  $P < 0.01$ ). Their movements were also related to the tide, since any sighted groups were 'down' the bay during flood tides ( $\chi^2$  test,  $P < 0.01$ ).

Surface feeding behaviours (actively chasing or circling fish, tossing a fish in the air and then retrieving it or holding prey in the mouth and aerial behaviours around the schools (*sensu* Lodi & Hetzel, 1999)) were documented in 15 encounters, including individual dolphins chasing and/or preying identified species by direct observation or photographs such as Lebranche mullet (*Mugil liza*) (Figure 3A), white mullet (*Mugil curema*) (Figure 3B), largehead hairtail (*Trichiurus lepturus*) and Brazilian menhaden (*Brevoortia aurea*). Interactions with other species included surface feeding of rough-toothed dolphins together with Brown boobies (*Sula leucogaster*) (in nine encounters), and the Magnificent frigatebird (*Fregata magnificens*) (in five encounters).

### Photo-identification

A total of 9402 photographs were taken, of which 3200 pictures (34%) were characterized as being of good or excellent quality ( $\geq 3$ ). In total 115 rough-toothed dolphins were individualized using photo-identification techniques, 61 (53%) of which were resighted on between one (47.5%) and four (9.8%) occasions, indicating some site fidelity (Supplementary Table S1). The interval between resightings ranged from 7 to 2087 days (mean = 268). Photo-identification resightings over one to six years indicated long-term site fidelity in coastal waters off Rio de Janeiro for at least some individuals.

The discovery curve continued to rise throughout the study period, with newly sighted individuals being identified (Figure 4).

Through the AHC analysis, 30 individuals (49.2%) were grouped in the low degree of fidelity, 12 (19.7%) in the medium




Fig. 3. Rough-toothed dolphins (*Steno bredanensis*) capturing Lebranche mullet (*Mugil liza*) (A) (photo: Liliane Lodi) and white mullet (*Mugil curema*) (B) (photo: Bia Hetzel) in Guanabara Bay.

degree and 19 (31.1%) in the high degree (Figure 5). The reference values found in each parameter and degree of fidelity are shown in Table 2. The resulting CCC was 0.76, indicating an adequate clustering of the data.

Some identified rough-toothed dolphins were sighted on the same dates and groups. Trios and pairs were sighted together three and four times respectively (Table 3).

### Discussion

The rough-toothed dolphin has been described as a primarily pelagic cetacean species that is found in tropical and subtropical oceans throughout the world; however, information on its distribution patterns or habitat use in most coastal regions remains limited. We present the first uninterrupted investigation of rough-toothed dolphins in south-eastern Brazil, indicating that in coastal environments this species may show peaks of occurrence in autumn and winter, and low site fidelity, despite inter-annual resightings. There is also indication of a relationship between the presence of groups near rocky shores during the closing season of mullet fisheries.

The peak occurrence of rough-toothed dolphins in this study corresponds to the reproductive or recruitment season of Lebranche mullet, an important item of their diet (Lodi & Hetzel, 1999). *Mugil* sp., *M. curema* and *T. lepturus* have also been previously described in the diet of *S. bredanensis* (Lodi & Hetzel, 1999; Di Benedetto *et al.*, 2001; Lodi *et al.*, 2012). Our study shows that preferred areas for rough-toothed dolphins consist of rocky coastlines, which may be found in the interior of Guanabara Bay, this may help them in pushing fish schools against the shoreline (Lodi & Hetzel, 1999) and coral reefs (Rossi-Santos *et al.*, 2006).

Within the Delphinidae family several species are known or suspected to make use of tidal features presumably because of




Fig. 4. Cumulative sighting and resighting frequency of rough-toothed dolphins (*Steno bredanensis*) along coastal waters off Rio de Janeiro from August 2011 to April 2018.

enhanced foraging opportunities (Benjamins *et al.*, 2015). Our results also show that groups usually moving ‘up’ Guanabara Bay would do so through areas of greater depths and moving ‘down’ through lower depths, closer to coastal areas. Groups were also mostly sighted ‘down’ the bay during flood tides. The counter-current movement (‘down’ movements during flood tide) may represent a foraging method. Dolphins may catch fish more easily when fish are swimming with or being carried by the current (Shane, 1980).

Group sizes were consistent with those previously reported for Brazil (1–50: Lodi & Hetzel, 1998b; Ramos *et al.*, 2010) and known patterns of social groupings for *S. bredanensis* in other regions, e.g. Mauritania (10–12: Addink & Smeenk, 2001), the Canary Islands (10–50: Ritter, 2002), French Polynesia (1–35: Gannier & West, 2005; 1–23: Oremus *et al.*, 2012), Honduras (5–30: Kuczaj & Yeater, 2007) and Hawaiian Archipelago (2–90: Baird *et al.*, 2008).


Groups with immature individuals were usually larger, offering more protection and possibly allowing better cognitive development (e.g. Hill *et al.*, 2007; Bender *et al.*, 2009; Kuczaj & Eskelinen, 2014; Mackey *et al.*, 2014). Rough-toothed dolphins appear to cooperate in a variety of ways to increase foraging success and may even actively teach immatures to forage (Steiner, 1995; Lodi & Hetzel, 1999; Addink & Smeenk, 2001; Pitman & Stinchcomb, 2002; Ritter, 2002).

Migratory movements of the species was first reported in Brazil by Lodi *et al.* (2012). Four photo-identified rough-toothed dolphins from Rio de Janeiro were resighted in the Cabo Frio region, ~117 and 119.7 km from the original sighting. Three individuals catalogued in this study were resighted in five different opportunities in the coastal waters between south and north of Rio de Janeiro, at intervals of one to five years, with a maximum linear distance of about 297 km (Wedekin *et al.*, 2017). Such resightings reinforce the hypothesis of degree of site fidelity. Moreover, three catalogued rough-toothed dolphins seen in Rio de Janeiro state were resighted 240 km southwards of São Paulo state (Santos *et al.*, 2019). The straight-line distance of 480 km between resighting locations in the main Hawaiian waters is the greatest travel distance reported for the species (Baird *et al.*, 2008), showing that *S. bredanensis* are

likely to have large home ranges. The shape of the discovery curve and the predominance of low site fidelity found for rough-toothed dolphins around inshore waters off Rio de Janeiro suggests a large home range and possibly a nomadic way of life.

Photo-identification studies of rough-toothed dolphins conducted in the Atlantic and Pacific archipelagos report long-term site fidelity in the species (i.e. individuals sighted over multiple years), suggesting the presence of resident populations near oceanic islands (Gannier & West, 2005; Mayr & Ritter, 2005; Kuczaj & Yeater, 2007; Baird *et al.*, 2008; Johnston *et al.*, 2008; Oremus *et al.*, 2012). Additionally, strong genetic differentiation among islands as well as among archipelagos of insular communities of rough-toothed dolphins was reported in the central Pacific Ocean from the Hawaiian, French Polynesian and Samoan archipelagos, where separate management units were identified (Albertson *et al.*, 2017). Site fidelity with little dispersal between populations has been shown, and rough-toothed dolphins can form highly localized, and apparently isolated, island communities, which is relevant for their conservation (Oremus *et al.*, 2012). Increased productivity near the Canary Islands, Hawaii and French Polynesia due to a variety of oceanographic processes, potentially results in better predictability of food resources in these ecosystems. Conversely, the discontinuity of food within the surrounding oligotrophic areas, might encourage high site fidelity and increased prey availability (Ritter, 2002; Gannier & West, 2005; Baird *et al.*, 2008; Oremus *et al.*, 2012; Albertson *et al.*, 2017).

Long-term and large-scale studies are needed to determine the range of home areas for groups of rough-toothed dolphins in south-eastern Brazil, as well as the ecological characteristics of their habitats. The social organization of rough-toothed dolphin groups is poorly understood; this species not only has strong social bonds between mother and calf/juvenile but, based on dolphin resightings, at least some animals in stable social groups for several years (Mayr & Ritter, 2005; Kuczaj & Yeater, 2007; Santos *et al.*, 2019). Data reported in this study on the associations between identified individuals support this hypothesis; thus, *S. bredanensis* exhibits behaviours associated with a complex social system such as formation of tight and synchronous



**Fig. 5.** Agglomerative hierarchical clustering (AHC) dendrogram of rough-toothed dolphins (*Steno bredanensis*) in coastal waters off Rio de Janeiro. The dashed lines represent clusters.

swimming, high tactile contact, cooperative foraging, and mass stranding events, provisioning of large prey to calves and care giving behaviours (Lodi & Hetzel, 1998a, 1999; Addink & Smeenk,

**Table 2.** Reference values found in each parameter (i, ii and iii) according to each degree of site fidelity of rough-toothed dolphins in coastal waters off Rio de Janeiro

Degrees	(i)		(ii)		(iii)	
	Min.	Max.	Min.	Max.	Min.	Max.
Low	0.12	0.75	0.09	0.14	0.25	0.50
Medium	0.60	1.00	0.09	0.19	0.25	0.50
High	0.22	0.50	0.14	0.23	0.50	1.00

**Table 3.** Individuals observed together in coastal waters off Rio de Janeiro. Number of sightings = times individuals were sighted. Number of associations = times individuals were sighted together

Individuals	Number of sightings	Number of associations
#030	5	4
#036	4	
#030	5	3
#036	4	
#071	3	
#048	4	3
#063	5	
#065	3	

2001; Pitman & Stinchcomb, 2002; Gotz *et al.*, 2006; Kuczaj & Yeater, 2007). These findings suggest social organization may play a role in rough-toothed dolphin population structure (Oremus *et al.*, 2012).

Guanabara Bay, an area used by rough-toothed dolphins, is one of the most industrialized coastal areas of Brazil (Fistarol *et al.*, 2015) harbouring industries, refineries, ports, shipyards and terminals of the oil and gas industries that contribute to water and noise pollution. Anthropogenic interference in Guanabara Bay area is high, with heavy eutrophication and hypoxia (Aguilar *et al.*, 2011), emergence of pathogenic microorganisms (Fistarol *et al.*, 2015), metal bioaccumulation in prey such as *M. liza* (Hauser-Davis *et al.*, 2016) and heavy shipping and vessel traffic (Bittencourt *et al.*, 2014).

In this study some records of rough-toothed dolphins were made in five gillnet fishing spots (or 'pesqueiros') of the artisanal fishing community of Copacabana, *Colônia de Pesca de Copacabana*, Z-13 (Amorim & Monteiro-Neto, 2016). Known prey of rough-toothed dolphins such as white mullet, largehead hairtail and Brazilian menhaden are fished in gillnets at these fishing spots (Amorim & Monteiro-Neto, 2016), although the potential threats of by-catch and associated dolphin mortality have yet to be reported. No sightings of *S. bredanensis* occurred in the MoNaCa Cagarras and adjacent areas but rather in Resex Itaipu where fishing activity occurs which again points to the need for further monitoring potential interactions with fisheries.

This study shows that at least some individuals of rough-toothed dolphins use coastal habitats off Rio de Janeiro, making them more exposed than their pelagic conspecifics to a variety of anthropogenic disturbances such as harassment from boats, risk of incidental capture, over-fishing, organic and persistent pollution and marine traffic. The long-term effects of anthropogenic activities on the survival of this species are still unknown. Nevertheless, sound management and continued monitoring efforts are essential to ensure the conservation of *S. bredanensis* along the coast of Rio de Janeiro. Our results may contribute

towards forming a benchmark for further studies in this region, which are fundamental to fill gaps in our understanding of this species throughout its distribution. Within the Atlantic Ocean basin three populations were detected (southern, south-eastern Brazil and Caribbean), and are considered as population management units for conservation purposes (Da Silva *et al.*, 2015).

Further data collection of environmental and anthropic variables is equally important for a better understanding of the factors driving rough-toothed dolphin habitat use and to allow future habitat modelling. Continuous photo-identification surveys, through enhanced research collaboration, are also of great importance to improve our understanding of the home range and population status of rough-toothed dolphins.

**Supplementary material.** The supplementary material for this article can be found at <https://doi.org/10.1017/S0025315420000132>.

**Acknowledgements.** We would like to thank Bruno F.T. Simões and Erick Monteiro for the support regarding the statistical tests used in this study and Monica Borobia, Eduardo Mortheo and an anonymous reviewer for improving our manuscript immensely with their comments.

**Financial support.** The results of this study are part of the Ilhas do Rio I, II and III Project, of the Mar Adentro Institute, which is sponsored by Petrobras (grant numbers 6000.0064815.11.2, 6000.0086840.13.2 and 5850.0106133.17.2, respectively). Baleias e Golfinhos do Rio de Janeiro Project was supported by Marine Program of WWF-Brazil (grant number CPT 00776-2016) and the SOS Mata Atlântica Foundation (unnumbered donation).

## References

- Addink MJ and Smeenk C (2001) Opportunistic feeding behaviour of rough-toothed dolphins *Steno bredanensis* off Mauritania. *Zoologische Verhandlungen* 334, 38–48.
- Aguiar VMC, Neto JAB and Rangel CM (2011) Eutrophication and hypoxia in four streams discharging in Guanabara Bay, RJ, Brazil, a case study. *Marine Pollution Bulletin* 62, 1915–1919.
- Albertson GR, Baird RW, Oremus M, Poole MM, Matien KK and Baker CS (2017) Staying close to home? Genetic differentiation of rough-toothed dolphins near oceanic islands in the central Pacific Ocean. *Conservation Genetics* 18, 33–51. doi: 10.1007/s10592-016-0880-z.
- Amorim RB and Monteiro-Neto C (2016) Marine protected area and the spatial distribution of the gill net fishery in Copacabana, Rio de Janeiro, RJ, Brazil. *Brazilian Journal of Biology* 76, 1–9. <http://dx.doi.org/10.1590/1519-6984.06614>.
- Baird RW, Webster DL, Mahaffy SD, MCSweeney DJ, Schorr GS and Ligon DA (2008) Site fidelity and association patterns in a deep-water dolphin: rough-toothed dolphins (*Steno bredanensis*) in the Hawaiian Archipelago. *Marine Mammal Science* 24, 535–553.
- Bender CE, Herzing DL and Bjorklund DF (2009) Evidence of teaching in Atlantic spotted dolphins (*Stenella frontalis*) by mother dolphins foraging in the presence of their calves. *Animal Cognition* 12, 43–53.
- Benjamins S, Dale A, Hastie G, Waggitt JJ, Lea M, Scott B and Wilson B (2015) Confusion reigns? A review of marine megafauna interactions with tidal-stream environments. *Oceanography and Marine Biology* 53, 1–54.
- Bittencourt L, Carvalho RR, Lailson-Brito J and Azevedo AF (2014) Underwater noise pollution in a coastal tropical environment. *Marine Pollution Bulletin* 83, 331–336.
- Choi L, Blume JD and Dupont WD (2015) Elucidating the foundations of statistical inference with  $2 \times 2$  tables. *PLoS ONE* 10, e0121263.
- Da Silva DMP, Azevedo AF, Secchi ER, Barbosa LA, Flores PAC, Carvalho RR, Bisi T, Lailson-Brito J and Cunha HA (2015) Molecular taxonomy and population structure of the rough-toothed dolphin *Steno bredanensis* (Cetartiodactyla: Delphinidae). *Zoological Journal of the Linnean Society* 175, 949–962.
- Day J, Dudley N, Hockings M, Holmes G, Laffoley DDA, Stolton S and Wells SM (2012) Guidelines for applying the IUCN protected area management categories to marine protected areas. *Best Practice Protected Area Guidelines Series No. 19*. Gland: IUCN.
- DeGroot MH and Schervish MJ (2012) *Probability and Statistics*, 4th Edn. Boston, MA: Pearson Education.
- Di Benedetto AP, Ramos RM and Lima NRW (1998) Fishing activity in northern Rio de Janeiro State (Brazil) and its relation with small cetaceans. *Brazilian Archives of Biology and Technology* 3, 1–7.
- Di Benedetto APM, Ramos RMA, Siciliano S, Santos RA, Bastos G and Fagundes-Neto E (2001) Stomach contents of delphinids from Rio de Janeiro, southeastern Brazil. *Aquatic Mammals* 27, 24–28.
- Dorneles PR, Lailson-Brito J, Santos RA, Costa PAS, Malm O, Azevedo AF and Torres JPM (2007) Cephalopods and cetaceans as indicators of offshore bioavailability of cadmium off Central South Brazil Bight. *Environmental Pollution* 148, 352–359.
- Espécie MA, Tardin RHO and Simão SM (2010) Degrees of residence of Guiana dolphins (*Sotalia guianensis*) in Ilha Grande Bay, south-eastern Brazil: a preliminary assessment. *Journal of the Marine Biological Association of the United Kingdom* 90, 1633–1639.
- Fistarol GO, Coutinho FH, Moreira APB, Venas T, Cánovas A, de Paula SE, Coutinho R, Mora RL, Valentin JL, Tenenbaum DR, Paranhos R, Valle RAB, Vicente ACP, Amado Filho GM, Pereira RC, Kruger R, Rezende CE, Thompson CC, Salomon P and Thompson FL (2015) Environmental and sanitary conditions of Guanabara Bay, Rio de Janeiro. *Frontiers in Microbiology* 6, 1232.
- Gannier A and West KL (2005) Distribution of the rough-toothed dolphin (*Steno bredanensis*) around the Windward Islands (French Polynesia). *Pacific Science* 59, 17–24.
- Gotz T, Verfass UK and Schnitzler HU (2006) Eavesdropping in wild rough-toothed dolphins (*Steno bredanensis*)? *Biology Letters* 2, 5–7.
- Greenacre M (1988) Correspondence analysis of multivariate categorical data by weighted least-squares. *Biometrika* 75, 457–467.
- Hauser-Davis RA, Isabella CAC, Bordon CA C, Oliveira TF and Zioli RL (2016) Metal bioaccumulation in edible target tissues of mullet (*Mugil liza*) from a tropical bay in southeastern Brazil. *Journal of Trace Elements in Medicine and Biology* 36, 38–43.
- Hill HM, Greer T, Solangi M and Kuczaj SA (2007) All mothers are not the same: maternal styles in bottlenose dolphins (*Tursiops truncatus*). *International Journal of Comparative Psychology* 20, 35–54.
- Instituto Brasileiro de Geografia e Estatística – IBGE (2018) Brasil – Rio de Janeiro. Available at <https://cidades.ibge.gov.br/brasil/rj/rio-de-janeiro/panorama> (Accessed 30 September 2018).
- Johnston DW, Robbins J, Chapla ME, Matthila DK and Andrews KR (2008) Diversity, habitat associations and stock structure of Odontocete cetaceans in the waters of American Samoa, 2003–06. *Journal of Cetacean Research and Management* 10, 59–66.
- Kjerfve B, Ribeiro CHA, Dias GTM, Filippo AM and Quaresma VS (1997) Oceanographic characteristics of an impacted coastal bay: Baía de Guanabara, Rio de Janeiro, Brazil. *Continental Shelf Research* 17, 1609–1643.
- Kuczaj SA and Yeater DB (2007) Observations of rough-toothed dolphins (*Steno bredanensis*) off the coast of Utila, Honduras. *Journal of the Marine Biological Association of the United Kingdom* 87, 141–148.
- Kuczaj SA and Eskelinen HC (2014) Why do dolphins play? *Animal Behavior and Cognition* 1, 113–127.
- Lailson-Brito J, Dorneles PR, Azevedo-Silva CE, Bisi TL, Vidal LG, Legat LN, Azevedo AF, Torres JPM and Malm O (2012) Organochlorine compound accumulation in delphinids from Rio de Janeiro State, southeastern Brazilian coast. *Science of the Total Environment* 433, 123–131.
- Legendre P and Legendre L (2012) *Numerical Ecology*, 3rd Edn. Amsterdam: Elsevier Science B.V.
- Lodi L and Capistrano L (1990) Capturas acidentais de pequenos cetáceos no litoral norte do Estado do Rio de Janeiro. *Biotemas* 3, 47–65.
- Lodi L and Hetzel B (1998a) Investigating a mass stranding of rough-toothed dolphins in Brazil. *Whalewatcher. Journal of the American Cetacean Society* 31, 18–20.
- Lodi L and Hetzel B (1998b) O golfinho-de-dentes-rugosos (*Steno Bredanensis*) no Brasil. *Bioikos* 12, 29–45.
- Lodi L and Hetzel B (1999) Rough-toothed dolphin, *Steno Bredanensis*, feeding behaviors in Ilha Grande Bay, Brazil. *Biociências* 7, 29–42.
- Lodi L and Tardin R (2018) Citizen science contributes to the understanding of the occurrence and distribution of cetaceans in southeastern Brazil – a case study. *Ocean and Coastal Management* 158, 45–55.
- Lodi L, Oliveira RHT, Figueiredo LD and Simão SM (2012) Movements of the rough-toothed dolphin (*Steno bredanensis*) in Rio de Janeiro State, south-eastern Brazil. *Marine Biodiversity Records* 5, 1–4.



- Mackey AD, Makecha RN and Kuczaj SA** (2014) The development of social play in bottlenose dolphins (*Tursiops truncatus*). *Animal Behavior and Cognition* **1**, 19–35.
- Mason RL, Gunst RF and Hess JL** (2003) *Statistical Design and Analysis of Experiments with Application to Engineering and Science*, 2nd Edn. Hoboken, NJ: John Wiley & Sons.
- Mayr I and Ritter F** (2005) Photo-identification of rough-toothed dolphins (*Steno bredanensis*) off La Gomera (Canary Islands) with new insights into social organization. In Evans PGH and Ridoux V (eds), *Proceedings of the Nineteenth Annual Conference of the European Cetacean Society Annual Meeting, L'Espace Encan Conference Centre, La Rochelle, 2–7 April 2005. Marine Mammals and Food: From Organisms to Ecosystems*. France: Series European Research on Cetaceans v. 19, pp. 1–3.
- Meirelles ACO and Barros HMDR** (2007) Plastic debris ingested by a rough-toothed dolphin, *Steno bredanensis*, stranded alive in northeastern Brazil. *Biotemas* **20**, 127–131.
- Monteiro-Neto C, Alves-Júnior TT, Ávila FJC, Campos AA, Costa AF, Silva CPN and Furtado-Neto MAA** (2000) Impact of fisheries on the tucuxi (*Sotalia fluviatilis*) and rough toothed dolphin (*Steno bredanensis*) populations off Ceará state, northeastern Brazil. *Aquatic Mammals* **26**, 49–56.
- Netto RF and Barbosa LA** (2003) Cetaceans and fishery interactions along the Espírito Santo State, southeastern Brazil during 1994–2001. *Latin American Journal of Aquatic Mammals* **2**, 57–60.
- Oremus M, Poole MM, Steel D and Baker SC** (2007) Isolation and inter-change among insular spinner dolphin communities in the South Pacific revealed by individual identification and genetic diversity. *Marine Ecology Progress Series* **336**, 275–289.
- Oremus M, Poole MM, Albertson GR and Baker SC** (2012) Pelagic or insular? Genetic differentiation of rough-toothed dolphins in the Society Islands, French Polynesia. *Journal of Experimental Marine Biology and Ecology* **432–433**, 37–46.
- Passadore C, Möller L, Diaz-Aguirre F and Parra GJ** (2018) High site fidelity and restricted ranging patterns in southern Australian bottlenose dolphins. *Ecology and Evolution* **8**, 242–256. doi: 10.1002/ece3.3674.
- Pitman RL and Stinchcomb C** (2002) Rough-toothed dolphins (*Steno bredanensis*) as predators of mahimahi (*Coryphaena hippurus*). *Pacific Science* **56**, 447–450.
- Ramos R, Poletto F, Moreira S, Erber C, Dafferner G, Freitas R, Figna V, Miranda C, Alencastro P, Carneiro A, Fortes R, Rinaldi G, Demari e Silva E and Barbosa M** (2010) Família Delphinidae: outros pequenos golfinhos. In Ramos RMA, Siciliano S and Ribeiro R (eds), *Monitoramento da biota marinha em navios de sismica: seis anos de pesquisa (2001–2007)*. Vitória, ES, Brazil: Everest Tecnologia em Serviços/PGS Investigação Petrolífera Ltda., pp. 649–728.
- Ritter F** (2002) Behavioral observations of rough-toothed dolphins (*Steno bredanensis*) off La Gomera, Canary Islands (1995–2000), with special reference to their interactions with humans. *Aquatic Mammals* **28**, 46–59.
- Rohlf FJ** (1970) Adaptive hierarchical clustering schemes. *Systematic Zoology* **19**, 58–82.
- Rossi-Santos MR, Wedekin L and Sousa-Lima RS** (2006) Distribution and habitat use of small cetaceans in the Abrolhos Bank, eastern Brazil. *Latin American Journal of Aquatic Mammals* **5**, 23–28.
- Ruellan F** (1944) A evolução geomorfológica da Baía de Guanabara e das regiões vizinhas. *Revista Brasileira de Geografia* **4**, 3–66.
- Santos MCO, Lailson-Brito J, Flach L, Oshima JEF, Figueiredo GC, Carvalho RR, Ventura ES, Molina JMB and Azevedo AF** (2019) Cetacean movements in coastal waters of the southwestern Atlantic ocean. *Biota Neotropica* **19**, e20180670. <http://dx.doi.org/10.1590/1676-0611-BN-2018-0670>.
- Shane S H** (1980) Occurrence, movements, and distribution of bottlenose dolphin, *Tursiops truncatus*, in southern Texas. *Fishery Bulletin* **83**, 593–601.
- Shane S, Wells RS and Würsig B** (1986) Ecology, behavior and social organization of the bottlenose dolphin: a review. *Marine Mammal Science* **2**, 34–63.
- Sokal RR and Rohlf FJ** (1962) The comparison of dendrograms by objective methods. *Taxon* **11**, 33–40.
- Steiner L** (1995) Rough-toothed dolphin, *Steno bredanensis*: a new species record for the Azores, with some notes on behaviour. *Arquipélago* **13A**, 125–127.
- van Weerelt M, Cunha L, Dorneles PR, Padilha, Ormond J, Torres F, Torres JP, Batista D, Nudi A, Wagener A, Cabral A, Pinto F and Paranhos R** (2013) Monitoramento das águas e dos poluentes no MoNa das Ilhas Cagarras e entorno. In Moraes F, Bertoncini A and Aguiar A (eds) *História, pesquisa e biodiversidade do Monumento Natural das Ilhas Cagarras*. Rio de Janeiro, RJ, Brazil, pp. 228–243. [Museu Nacional Série Livros, no. 48].
- Wedekin LL, Rossi-Santos MR, Baracho C, Cypriano-Souza AL and Simões-Lopes PC** (2014) Cetacean records along a coastal-offshore gradient in the Vitória-Trindade Chain, western South Atlantic Ocean. *Brazilian Journal of Biology* **74**, 137–144.
- Wedekin L, Paro A, Cypriano A, Daura-Jorge F, Silveira F, Olimpio J, Dalpaz L, Rossi-Santos M and Cremer M** (2017) PMC-BS/Projeto de Monitoramento de Cetáceos da Baía de Santos. *Segundo Relatório Anual, Ciclos 1 a 4*. Socioambiental Consultores Associados, 815 pp.
- White GC and Garrot RA** (1990) *Analysis of Wildlife Radio-Tracking Data*. San Diego, CA: Academic Press.
- Whitehead H, Christal J and Tyack, PL** (2000) Studying cetacean social structure in space and time: Innovative. In Mann J, Connor RC, Tyack PL and Whitehead H (eds), *Cetacean Societies*. Chicago, IL: University of Chicago Press, pp. 65–87.
- Würsig B and Jefferson TA** (1990) Methodology of photo-identification for small cetaceans. In Hammond PS, Mizroch SA and Donovan GP (eds), *Individual Recognition of Cetaceans: Use of Photoidentification and Other Techniques to Estimate Population Parameters*. Cambridge: International Whaling Commission, Special Issue 12, pp. 43–52.
- Yogui GT, Santos MCO, Bertozzi CP and Montone RC** (2010) Levels of persistent organic pollutants and residual pattern of DDTs in small cetaceans from the coast of São Paulo, Brazil. *Marine Pollution Bulletin* **60**, 1862–1867.